

Care Quality Commission

Inspection Evidence Table

St Melor House Surgery (1-551300146)

Inspection date: **14 August 2018**

Date of data download: 08 August 2018

Please note: Any Quality Outcomes Framework (QOF) data relates to 2016/17.

Safe

Safety systems and processes

Safeguarding	Y/N
There was a lead member(s) of staff for safeguarding processes and procedures.	Y
Safety and safeguarding systems, processes and practices were developed, implemented and communicated to staff.	Y
Policies were in place covering adult and child safeguarding.	Y
Policies were updated and reviewed and accessible to all staff.	Y
Partners and staff were trained to appropriate levels for their role (for example level three for GPs, including locum GPs)	Y

Effective

Effective needs assessment, care and treatment

Prescribing				
Indicator	Practice performance	CCG average	England average	England comparison
Average daily quantity of Hypnotics prescribed per Specific Therapeutic group Age-sex Related Prescribing Unit (STAR PU) (01/04/2017 to 31/03/2018) (NHSBSA)	0.85	0.82	0.84	Comparable with other practices

People with long-term conditions

Diabetes Indicators				
Indicator	Practice performance	CCG average	England average	England comparison
The percentage of patients with diabetes, on the register, in whom the last IFCC-HbA1c is 64 mmol/mol or less in the preceding 12 months (01/04/2016 to 31/03/2017) (QOF)	76.7%	83.5%	79.5%	Comparable with other practices
QOF Exceptions	Practice Exception rate (number of exceptions) 8.9% (25)	CCG Exception rate 18.1%	England Exception rate 12.4%	
Indicator	Practice performance	CCG average	England average	England comparison
The percentage of patients with diabetes, on the register, in whom the last blood pressure reading (measured in the preceding 12 months) is 140/80 mmHg or less (01/04/2016 to 31/03/2017) (QOF)	81.8%	79.6%	78.1%	Comparable with other practices
QOF Exceptions	Practice Exception rate (number of exceptions) 6.4% (18)	CCG Exception rate 12.2%	England Exception rate 9.3%	

Indicator	Practice performance	CCG average	England average	England comparison
The percentage of patients with diabetes, on the register, whose last measured total cholesterol (measured within the preceding 12 months) is 5 mmol/l or less (01/04/2016 to 31/03/2017) (QOF)	85.6%	82.7%	80.1%	Comparable with other practices
QOF Exceptions	Practice Exception rate (number of exceptions)	CCG Exception rate	England Exception rate	
	6.4% (18)	17.6%	13.3%	

Other long term conditions				
Indicator	Practice	CCG average	England average	England comparison
The percentage of patients with asthma, on the register, who have had an asthma review in the preceding 12 months that includes an assessment of asthma control using the 3 RCP questions, NICE 2011 menu ID: NM23 (01/04/2016 to 31/03/2017) (QOF)	71.6%	78.1%	76.4%	Comparable with other practices
QOF Exceptions	Practice Exception rate (number of exceptions)	CCG Exception rate	England Exception rate	
	2.0% (8)	9.6%	7.7%	
Indicator	Practice	CCG average	England average	England comparison
The percentage of patients with COPD who have had a review, undertaken by a healthcare professional, including an assessment of breathlessness using the Medical Research Council dyspnoea scale in the preceding 12 months (01/04/2016 to 31/03/2017) (QOF)	92.5%	92.7%	90.4%	Comparable with other practices
QOF Exceptions	Practice Exception rate (number of exceptions)	CCG Exception rate	England Exception rate	
	3.6% (4)	13.6%	11.4%	

Indicator	Practice	CCG average	England average	England comparison
The percentage of patients with hypertension in whom the last blood pressure reading measured in the preceding 12 months is 150/90mmHg or less (01/04/2016 to 31/03/2017) (QOF)	87.9%	83.9%	83.4%	Comparable with other practices

QOF Exceptions	Practice Exception rate (number of exceptions)	CCG Exception rate	England Exception rate	
	2.8% (23)	4.5%	4.0%	
Indicator	Practice	CCG average	England average	England comparison
In those patients with atrial fibrillation with a record of a CHA2DS2-VASc score of 2 or more, the percentage of patients who are currently treated with anti-coagulation drug therapy (01/04/2016 to 31/03/2017) (QOF)	86.1%	90.5%	88.4%	Comparable with other practices
QOF Exceptions	Practice Exception rate (number of exceptions)	CCG Exception rate	England Exception rate	
	12.2% (11)	8.3%	8.2%	
Any additional evidence or comments				

Families, children and young people

Child Immunisation				
Indicator	Numerator	Denominator	Practice %	Comparison to WHO target
The percentage of children aged 1 who have completed a primary course of immunisation for Diphtheria, Tetanus, Polio, Pertussis, Haemophilus influenza type b (Hib)(i.e. three doses of DTaP/IPV/Hib) (01/04/2016 to 31/03/2017)(NHS England)	69	71	97.2%	Met 95% WHO based target (significant variation positive)
The percentage of children aged 2 who have received their booster immunisation for Pneumococcal infection (i.e. received Pneumococcal booster) (PCV booster) (01/04/2016 to 31/03/2017) (NHS England)	62	67	92.5%	Met 90% minimum (no variation)
The percentage of children aged 2 who have received their immunisation for Haemophilus influenza type b (Hib) and Meningitis C (MenC) (i.e. received Hib/MenC booster) (01/04/2016 to 31/03/2017) (NHS England)	62	67	92.5%	Met 90% minimum (no variation)
The percentage of children aged 2 who have received immunisation for measles, mumps and rubella (one dose of MMR) (01/04/2016 to 31/03/2017) (NHS England)	62	67	92.5%	Met 90% minimum (no variation)

Working age people (including those recently retired and students)

Cancer Indicators				
Indicator	Practice	CCG average	England average	England comparison
The percentage of women eligible for cervical cancer screening at a given point in time who were screened adequately within a specified period (within 3.5 years for women aged 25 to 49, and within 5.5 years for women aged 50 to 64) (01/04/2016 to 31/03/2017) (Public Health England)	79.2%	75.9%	72.1%	Comparable with other practices
Females, 50-70, screened for breast cancer in last 36 months (3 year coverage, %) (PHE)	75.7%	76.3%	70.3%	N/A
Persons, 60-69, screened for bowel cancer in last 30 months (2.5 year coverage, %)(PHE)	59.4%	62.9%	54.5%	N/A
The percentage of patients with cancer, diagnosed within the preceding 15 months, who have a patient review recorded as occurring within 6 months of the date of diagnosis. (PHE)	71.9%	66.0%	71.2%	N/A
Number of new cancer cases treated (Detection rate: % of which resulted from a two week wait (TWW) referral) (01/04/2016 to 31/03/2017) (PHE)	45.7%	46.5%	51.6%	Comparable with other practices

People experiencing poor mental health (including people with dementia)

Mental Health Indicators				
Indicator	Practice	CCG average	England average	England comparison
The percentage of patients with schizophrenia, bipolar affective disorder and other psychoses who have a comprehensive, agreed care plan documented in the record, in the preceding 12 months (01/04/2016 to 31/03/2017) (QOF)	92.9%	94.0%	90.3%	Comparable with other practices
QOF Exceptions	Practice Exception rate (number of exceptions)	CCG Exception rate	England Exception rate	
	6.7% (1)	15.1%	12.5%	
Indicator	Practice	CCG average	England average	England comparison
The percentage of patients with schizophrenia, bipolar affective disorder and other psychoses whose alcohol consumption has been recorded in the preceding 12 months (01/04/2016 to 31/03/2017) (QOF)	92.3%	93.3%	90.7%	Comparable with other practices
QOF Exceptions	Practice Exception rate (number of exceptions)	CCG Exception rate	England Exception rate	
	13.3% (2)	13.8%	10.3%	

Indicator	Practice	CCG average	England average	England comparison
The percentage of patients diagnosed with dementia whose care plan has been reviewed in a face-to-face review in the preceding 12 months (01/04/2016 to 31/03/2017) (QOF)	91.1%	87.2%	83.7%	Comparable with other practices
QOF Exceptions	Practice Exception rate (number of exceptions)	CCG Exception rate	England Exception rate	
	6.7% (4)	8.6%	6.8%	

Monitoring care and treatment

Indicator	Practice	CCG average	England average
Overall QOF score (out of maximum 559)	553	553	539
Overall QOF exception reporting (all domains)	3.9%	6.2%	5.7%

Helping patients to live healthier lives

Indicator	Practice	CCG average	England average	England comparison
The percentage of patients with any or any combination of the following conditions: CHD, PAD, stroke or TIA, hypertension, diabetes, COPD, CKD, asthma, schizophrenia, bipolar affective disorder or other psychoses whose notes record smoking status in the preceding 12 months (01/04/2016 to 31/03/2017) (QOF)	97.6%	95.6%	95.3%	Comparable with other practices
QOF Exceptions	Practice Exception rate (number of exceptions)	CCG Exception rate	England Exception rate	
	0.6% (8)	0.8%	0.8%	

Well-led

Governance arrangements

Examples of structures, processes and systems in place to support the delivery of good quality and sustainable care.	
Practice specific policies	The practice had monitored their training requirements to ensure these were up to date. All relevant staff had completed Mental Capacity Act (MCA) training. We saw documentary evidence that the practice had a training matrix which scheduled mandatory and other training when due for renewal.

Feedback from patients
We spoke to three patients who were all happy with the service received and felt fully involved in decisions about their care and treatment.

Any additional evidence
The practice is currently collating and analysing its patient responses from a recent (July – August 2018) survey.

DO NOT DELETE THE NOTES BELOW

Notes: CQC GP Insight

GP Insight assesses a practice's data against all the other practices in England. We assess relative performance for the majority of indicators using a "z-score" (this tells us the number of standard deviations from the mean the data point is), giving us a statistical measurement of a practice's performance in relation to the England average. We highlight practices which significantly vary from the England average (in either a positive or negative direction). We consider that z-scores which are higher than +2 or lower than -2 are at significant levels, warranting further enquiry. Using this technique we can be 95% confident that the practice's performance is genuinely different from the average. It is important to note that a number of factors can affect the Z score for a practice, for example a small denominator or the distribution of the data. This means that there will be cases where a practice's data looks quite different to the average, but still shows as comparable, as we do not have enough confidence that the difference is genuine. There may also be cases where a practice's data looks similar across two indicators, but they are in different variation bands.

The percentage of practices which show variation depends on the distribution of the data for each indicator, but is typically around 10-15% of practices. The practices which are not showing significant statistical variation are labelled as comparable to other practices.

N.B. Not all indicators in the evidence table are part of the GP insight set and those that aren't will not have a variation band.

The following language is used for showing variation:

	Variation Band	Z-score threshold
1	Significant variation (positive)	$Z \leq -3$
2	Variation (positive)	$-3 < Z \leq -2$
3	Comparable to other practices	$-2 < Z < 2$
4	Variation (negative)	$2 \leq Z < 3$
5	Significant variation (negative)	$Z \geq 3$
6	No data	Null

Note: for the following indicators the variation bands are different:

- Child Immunisation indicators. These are scored against the World Health Organisation target of 95% rather than the England average.

It is important to note that z-scores are not a judgement in themselves, but will prompt further enquiry, as part of our ongoing monitoring of GP practices.

Guidance and Frequently Asked Questions on GP Insight can be found on the following link:

<http://www.cqc.org.uk/what-we-do/how-we-use-information/monitoring-gp-practices>

Glossary of terms used in the data.

- **COPD:** Chronic Obstructive Pulmonary Disease
- **PHE:** Public Health England
- **QOF:** Quality and Outcomes Framework (see <https://qof.digital.nhs.uk>).
- **RCP:** Royal College of Physicians.
- **STAR-PU:** Specific Therapeutic Group Age-sex weightings Related Prescribing Units. These weighting allow more accurate and meaningful comparisons within a specific therapeutic group by taking into account the types of people who will be receiving that treatment. ([See NHS Choices for more details](#)).