

Care Quality Commission

Inspection Evidence Table

Woodside Surgery (1-541134350)

Inspection date: 20 June 2019

Date of data download: 19 June 2019

Overall rating: Good

Please note: Any Quality Outcomes Framework (QOF) data relates to 2017/18.

Effective

Rating: Good

Effective needs assessment, care and treatment

Patients' needs were assessed, and care and treatment was delivered in line with current legislation, standards and evidence-based guidance supported by clear pathways and tools.

On our previous visit in October 2018 we rated the practice as requires improvement for providing effective services.

This was because there were significant gaps in relation to mandatory training uptake; notably infection prevention and control and fire safety. In addition, induction processes were not consistently applied and staff appraisals were out of date. Furthermore, processes for dissemination of clinical updates and alerts were not sufficiently clear.

At this inspection, we found the provider had satisfactorily addressed these issues and we have rated them as good for providing effective services.

	Y/N/Partial
The practice had systems and processes to keep clinicians up to date with current evidence-based practice.	Y
Patients' immediate and ongoing needs were fully assessed. This included their clinical needs and their mental and physical wellbeing.	Y
We saw no evidence of discrimination when staff made care and treatment decisions.	Y
Patients' treatment was regularly reviewed and updated.	Y
There were appropriate referral pathways were in place to make sure that patients' needs were addressed.	Y
Patients were told when they needed to seek further help and what to do if their condition deteriorated.	Y
Explanation of any answers and additional evidence: The provider had overhauled their approach to management of clinical and other patient safety alerts. A new protocol had been developed, which provided for a number of key personnel to receive and disseminate the alerts. Clinical alerts were discussed as a standing agenda item at clinical meetings, with a system for staff not present at meetings to sign to indicate they had been informed of any changes. A spreadsheet had been developed demonstrating when actions were required and completed.	

Prescribing	Practice performance	CCG average	England average	England comparison
Average daily quantity of Hypnotics prescribed per Specific Therapeutic group Age-sex Related Prescribing Unit (STAR PU) (01/01/2018 to 31/12/2018) <small>(NHSBSA)</small>	0.96	0.75	0.79	No statistical variation

Older people

Population group rating: Good

Findings
<ul style="list-style-type: none"> Older patients who were frail or may be vulnerable received a full assessment of their physical, mental and social needs. The practice used an appropriate tool to identify patients aged 65 and over who were living with moderate or severe frailty. Those identified as being frail had a clinical review including a review of medication. <ul style="list-style-type: none"> The practice followed up on older patients discharged from hospital. It ensured that their care plans and prescriptions were updated to reflect any extra or changed needs. Staff had appropriate knowledge of treating older people including their psychological, mental and communication needs. Patients over the age of 75 years were offered an annual holistic health review.

People with long-term conditions

Population group rating: **Good**

Findings

- One member of staff had recently qualified as a nurse associate. At the time of our visit the practice was developing a new approach to management of long-term conditions, utilising the skills of the nurse associate, to review multiple conditions at one appointment when possible, in the month of their birthday. Where patients' needs were more complex, practice staff co-ordinated care planning with other relevant health and care professionals.
- Staff who were responsible for reviews of patients with long-term conditions had received specific training.
- GPs followed up patients who had received treatment in hospital or through out of hours services for an acute exacerbation of asthma.
- The practice had recently introduced a system of same day appointment availability for practice nurse availability to deal with acute exacerbations of long-term conditions, such as asthma and COPD.

Diabetes Indicators	Practice	CCG average	England average	England comparison
The percentage of patients with diabetes, on the register, in whom the last IFCC-HbA1c is 64 mmol/mol or less in the preceding 12 months (01/04/2017 to 31/03/2018) <small>(QOF)</small>	76.3%	76.5%	78.8%	No statistical variation
Exception rate (number of exceptions).	15.1% (67)	8.2%	13.2%	N/A
The percentage of patients with diabetes, on the register, in whom the last blood pressure reading (measured in the preceding 12 months) is 140/80 mmHg or less (01/04/2017 to 31/03/2018) <small>(QOF)</small>	73.7%	74.7%	77.7%	No statistical variation
Exception rate (number of exceptions).	7.4% (33)	7.2%	9.8%	N/A

	Practice	CCG average	England average	England comparison
The percentage of patients with diabetes, on the register, whose last measured total cholesterol (measured within the preceding 12 months) is 5 mmol/l or less (01/04/2017 to 31/03/2018) <small>(QOF)</small>	79.9%	77.7%	80.1%	No statistical variation
Exception rate (number of exceptions).	15.8% (70)	11.8%	13.5%	N/A

Other long-term conditions	Practice	CCG average	England average	England comparison
The percentage of patients with asthma, on the register, who have had an asthma review in the preceding 12 months that includes an assessment of asthma control using the 3 RCP questions, NICE 2011 menu ID: NM23 (01/04/2017 to 31/03/2018) <small>(QOF)</small>	69.4%	73.8%	76.0%	No statistical variation
Exception rate (number of exceptions).	1.6% (11)	5.5%	7.7%	N/A
The percentage of patients with COPD who have had a review, undertaken by a healthcare professional, including an assessment of breathlessness using the Medical Research Council dyspnoea scale in the preceding 12 months (01/04/2017 to 31/03/2018) <small>(QOF)</small>	86.1%	85.7%	89.7%	No statistical variation
Exception rate (number of exceptions).	10.0% (28)	9.6%	11.5%	N/A

Indicator	Practice	CCG average	England average	England comparison
The percentage of patients with hypertension in whom the last blood pressure reading measured in the preceding 12 months is 150/90mmHg or less (01/04/2017 to 31/03/2018) (QOF)	80.9%	83.2%	82.6%	No statistical variation
Exception rate (number of exceptions).	2.0% (23)	3.1%	4.2%	N/A
In those patients with atrial fibrillation with a record of a CHA2DS2-VASc score of 2 or more, the percentage of patients who are currently treated with anti-coagulation drug therapy (01/04/2017 to 31/03/2018) (QOF)	85.6%	88.0%	90.0%	No statistical variation
Exception rate (number of exceptions).	3.2% (3)	6.0%	6.7%	N/A

Families, children and young people

Population group rating: Good

Findings

- Childhood immunisation uptake rates were in line with the World Health Organisation (WHO) targets.
- The practice had arrangements to identify and review the treatment of newly pregnant women on long-term medicines. These patients were provided with advice and post-natal support in accordance with best practice guidance.
- The practice had arrangements for following up failed attendance of children's appointments following an appointment in secondary care or for immunisation and would liaise with health visitors when necessary.
- Arrangements were in place for liaising with health visitors and other community staff, via the shared clinical system, for children and families with additional needs.

Child Immunisation	Numerator	Denominator	Practice %	Comparison to WHO target
The percentage of children aged 1 who have completed a primary course of immunisation for Diphtheria, Tetanus, Polio, Pertussis, Haemophilus influenza type b (Hib)((i.e. three doses of DTaP/IPV/Hib) (01/04/2017 to 31/03/2018) (NHS England)	116	119	97.5%	Met 95% WHO based target (significant variation positive)
The percentage of children aged 2 who have received their booster immunisation for Pneumococcal infection (i.e. received Pneumococcal booster) (PCV booster) (01/04/2017 to 31/03/2018) (NHS England)	109	115	94.8%	Met 90% minimum (no variation)
The percentage of children aged 2 who have received their immunisation for Haemophilus influenza type b (Hib) and Meningitis C (MenC) (i.e. received Hib/MenC booster) (01/04/2017 to 31/03/2018) (NHS England)	109	115	94.8%	Met 90% minimum (no variation)
The percentage of children aged 2 who have received immunisation for measles, mumps and rubella (one dose of MMR) (01/04/2017 to 31/03/2018) (NHS England)	108	115	93.9%	Met 90% minimum (no variation)

Working age people (including those recently retired and students)

Population group rating: Good

Findings

- Uptake of cervical screening was just under 72% which was in line with the national average. There were plans for the newly appointed nurse associate to receive training in cervical screening to provide additional resource to help improve access to appointments and improve uptake.
- The practice had systems to inform eligible patients to have the meningitis vaccine, for example before attending university for the first time.
- Patients had access to appropriate health assessments and checks including NHS checks for patients aged 40 to 74. There was appropriate and timely follow-up on the outcome of health assessments and checks where abnormalities or risk factors were identified.
- Patients could book or cancel appointments online and order repeat medication without the need to attend the surgery. At the time of our visit 14% of patients were registered for this service. The practice was promoting uptake, via their social media presence, to help them reach the target of

25% uptake.

Cancer Indicators	Practice	CCG average	England average	England comparison
The percentage of women eligible for cervical cancer screening at a given point in time who were screened adequately within a specified period (within 3.5 years for women aged 25 to 49, and within 5.5 years for women aged 50 to 64) (01/04/2017 to 31/03/2018) (Public Health England)	71.7%	76.3%	71.7%	No statistical variation
Females, 50-70, screened for breast cancer in last 36 months (3 year coverage, %) (01/04/2017 to 31/03/2018) (PHE)	62.9%	66.9%	70.0%	N/A
Persons, 60-69, screened for bowel cancer in last 30 months (2.5 year coverage, %)(01/04/2017 to 31/03/2018) (PHE)	53.1%	57.6%	54.5%	N/A
The percentage of patients with cancer, diagnosed within the preceding 15 months, who have a patient review recorded as occurring within 6 months of the date of diagnosis. (01/04/2017 to 31/03/2018) (PHE)	64.1%	65.2%	70.2%	N/A
Number of new cancer cases treated (Detection rate: % of which resulted from a two week wait (TWW) referral) (01/04/2017 to 31/03/2018) (PHE)	50.0%	49.1%	51.9%	No statistical variation

People whose circumstances make them vulnerable

Population group rating: Good

Findings

- End of life care was delivered in a coordinated way which took into account the needs of those whose circumstances may make them vulnerable.
- The practice held a register of patients living in vulnerable circumstances including those with a learning disability.
- The practice had a system for vaccinating patients with an underlying medical condition according to the recommended schedule.
- The practice was working within Calderdale Group Practice, and as part of a Primary Care Network to explore social prescribing options to offer additional support to people experiencing social isolation or loneliness.

**People experiencing poor mental health
(including people with dementia)**

Population group rating: Good

Findings

- The practice assessed and monitored the physical health of people with mental illness, severe mental illness, and personality disorder by providing access to health checks, interventions for physical activity, obesity, diabetes, heart disease, cancer and access to smoking cessation services.
- There was a system for following up patients who failed to attend for administration of long-term medication.
- When patients were assessed to be at risk of suicide or self-harm the practice had arrangements in place to help them to remain safe.
- Patients at risk of dementia were identified and offered an assessment to detect possible signs of dementia. When dementia was suspected there was an appropriate referral for diagnosis.
- The practice made use of community 'Quest' matrons who provided oversight to those people residing in care homes who were living with dementia.

Mental Health Indicators	Practice	CCG average	England average	England comparison
The percentage of patients with schizophrenia, bipolar affective disorder and other psychoses who have a comprehensive, agreed care plan documented in the record, in the preceding 12 months (01/04/2017 to 31/03/2018) (QOF)	91.8%	88.8%	89.5%	No statistical variation
Exception rate (number of exceptions).	7.6% (6)	12.9%	12.7%	N/A
The percentage of patients with schizophrenia, bipolar affective disorder and other psychoses whose alcohol consumption has been recorded in the preceding 12 months (01/04/2017 to 31/03/2018) (QOF)	83.3%	88.4%	90.0%	No statistical variation
Exception rate (number of exceptions).	8.9% (7)	11.9%	10.5%	N/A
The percentage of patients diagnosed with dementia whose care plan has been reviewed in a face-to-face review in the preceding 12 months (01/04/2017 to 31/03/2018) (QOF)	78.8%	80.2%	83.0%	No statistical variation
Exception rate (number of exceptions).	5.7% (2)	6.7%	6.6%	N/A

--

Effective staffing

The practice was able to demonstrate that staff had the skills, knowledge and experience to carry out their roles.

	Y/N/Partial
Staff had the skills, knowledge and experience to deliver effective care, support and treatment. This included specific training for nurses on immunisation and on sample taking for the cervical screening programme.	Y
The learning and development needs of staff were assessed.	Y
The practice had a programme of learning and development.	Y
Staff had protected time for learning and development.	Y
There was an induction programme for new staff.	Y
Induction included completion of the Care Certificate for Health Care Assistants employed since April 2015.	Y
Staff had access to regular appraisals, one to ones, coaching and mentoring, clinical supervision and revalidation. They were supported to meet the requirements of professional revalidation.	Y
The practice could demonstrate how they assured the competence of staff employed in advanced clinical practice, for example, nurses, paramedics, pharmacists and physician associates.	Y
There was a clear and appropriate approach for supporting and managing staff when their performance was poor or variable.	Y
<p>Explanation of any answers and additional evidence:</p> <p>The provider had introduced new systems of monitoring uptake of training, mandatory and otherwise for staff. Members of the management team had specific responsibility for oversight of this. Staff were provided with dedicated time to undertake necessary training. Where staff carried out training in their own time they were able to take time back or receive payment for additional hours worked. We saw an up to date staff training matrix which indicated that staff were up to date with regular training in line with the role they carried out in the practice. In addition, we heard of examples where staff had been encouraged to develop in their role and enhance their skills.</p>	

Notes: CQC GP Insight

GP Insight assesses a practice's data against all the other practices in England. We assess relative performance for the majority of indicators using a "z-score" (this tells us the number of standard deviations from the mean the data point is), giving us a statistical measurement of a practice's performance in relation to the England average. We highlight practices which significantly vary from the England average (in either a positive or negative direction). We consider that z-scores which are higher than +2 or lower than -2 are at significant levels, warranting further enquiry. Using this technique we can be 95% confident that the practice's performance is genuinely different from the average. It is important to note that a number of factors can affect the Z score for a practice, for example a small denominator or the distribution of the data. This means that there will be cases where a practice's data looks quite different to the average, but still shows as no statistical variation, as we do not have enough confidence that the difference is genuine. There may also be cases where a practice's data looks similar across two indicators, but they are in different variation bands.

The percentage of practices which show variation depends on the distribution of the data for each indicator, but is typically around 10-15% of practices. The practices which are not showing significant statistical variation are labelled as no statistical variation to other practices.

N.B. Not all indicators in the evidence table are part of the GP insight set and those that aren't will not have a variation band.

The following language is used for showing variation:

Variation Bands	Z-score threshold
Significant variation (positive)	≤ -3
Variation (positive)	> -3 and ≤ -2
Tending towards variation (positive)	> -2 and ≤ -1.5
No statistical variation	< 1.5 and > -1.5
Tending towards variation (negative)	≥ 1.5 and < 2
Variation (negative)	≥ 2 and < 3
Significant variation (negative)	≥ 3

Note: for the following indicators the variation bands are different:

- Child Immunisation indicators. These are scored against the World Health Organisation target of 95% rather than the England average.
- The percentage of respondents to the GP patient survey who responded positively to how easy it was to get through to someone at their GP practice on the phone uses a rules based approach for scoring, due to the distribution of the data. This indicator does not have a CCG average.

It is important to note that z-scores are not a judgement in themselves, but will prompt further enquiry, as part of our ongoing monitoring of GP practices.

Guidance and Frequently Asked Questions on GP Insight can be found on the following link:
<https://www.cqc.org.uk/guidance-providers/gps/how-we-monitor-gp-practices>

Glossary of terms used in the data.

- **COPD:** Chronic Obstructive Pulmonary Disease
- **PHE:** Public Health England
- **QOF:** Quality and Outcomes Framework
- **STAR-PU:** Specific Therapeutic Group Age-sex weightings Related Prescribing Units. These weighting allow more accurate and meaningful comparisons within a specific therapeutic group by taking into account the types of people who will be receiving that treatment.